

ON ADS_3 STRING THEORY

P. Marios Petropoulos

Centre de Physique Théorique

Ecole Polytechnique

F-91128 Palaiseau CEDEX

marios@cpht.polytechnique.fr

Abstract String theory on curved backgrounds has received much attention on account of both its own interest, and of its relation with gauge theories. Despite the progress made in various directions, several quite elementary questions remain unanswered, in particular in the very simple case of three-dimensional anti-de Sitter space. I briefly review these problems.

String theory is the most appropriate setting for studying quantum-gravity phenomena. In the absence of a truly non-perturbative approach, the usual method consists in analysing the string propagating on non-trivial backgrounds, that satisfy the requirements of conformal invariance of the corresponding sigma model.

Three-dimensional anti-de Sitter space was recognized long ago as a case of interest [1]–[5]. It is a maximally symmetric solution of Einstein’s equations with negative cosmological constant, and time is embedded non-trivially in the curved geometry. Alternatively, it corresponds to the Freedman–Gibbons electrovac solution of gauged supergravity, which can be shown to leave space-time supersymmetry unbroken [6]. Other peculiar features of ADS_3 are the absence of asymptotically flat regions, the presence of boundaries, as well as a rich causal structure, which makes it possible to obtain three-dimensional black holes after modding out some discrete symmetry [7].

As far as string theory is concerned, ADS_3 is an *exact* background, provided an NS–NS two-form is switched on. The underlying two-dimensional theory is a Wess–Zumino–Witten model on the $SU(1,1)$

*CPHT-PC-735.0999/September 1999

group manifold. Three-dimensional anti-de Sitter space is *the only* exact background – with a single time direction – where string propagation leads to a WZW model. This property does not hold for more general ADS_n .

Despite many efforts, driven in particular by the recent activity on ADS/CFT correspondence, and the apparent simplicity of the model at hand, several important and elementary issues are still beyond our understanding.

The analysis of string theory on ADS_3 plus torsion background can be performed in two steps. We first study the sigma model whose target space has the above geometry. Then, the latter has to be coupled to two-dimensional gravity. At the level of the Hilbert space, this amounts to the decoupling of a certain subspace, which becomes unphysical. Several questions must then be answered, among which: Is the physical spectrum free of negative-norm states? Is the operator algebra well-behaved, and what are the vertex operators and their interpretation? What is the one-loop modular-invariant partition function?

Very little is known about WZW models on non-compact groups, at a sufficiently rigorous and general level. Most of our knowledge is based on a formal extension of the compact case to some specific situations, and in the framework of current-algebra techniques. Target-space boundary conditions, in particular, are treated somehow carelessly, although we know how important they are for selecting various representations when studying quantum mechanics on ADS_3 [8].

We usually assume that the $SO(2,2) \cong SU(1,1)_L \times SU(1,1)_R$ symmetry of the above model is realized in terms of an affine Lie algebra, the level of which is not quantized because $\pi_3(SU(1,1)) = 0$. The world-sheet energy-momentum tensor is given by the affine Sugawara construction, and its modes satisfy the Virasoro algebra with central charge $c = \frac{3k}{k+2}$. Choosing $k < -2$ leads to one time and two space directions, while the central charge is positive.

Finally, the Hilbert space is formally constructed as in the compact case: it is a direct sum of products of representations of the left and right current algebras. Highest-weight representations of the $SU(1,1)$ current algebra are labelled by the spin j of the primary fields (states of level zero), which form a representation of the global algebra. Higher-level states are obtained by acting with negative-frequency current oscillators on the above primary states, which are annihilated by all positive-frequency modes.

Irreducible representations of the global algebra are of two kinds: discrete or continuous. The discrete ones have highest or lowest weight, whereas the continuous ones do not. They are all infinite-dimensional,

at least when unitarity is demanded. This is due to the non-compact nature of $SU(1,1)$, which also implies that the metric on the group manifold is not positive-definite.

Representations of the $SU(1,1)$ current algebra have always infinite towers of negative-norm states. Therefore the corresponding WZW model cannot be unitary. Nevertheless, since we are interested in the string theory, the relevant question is whether the physical spectrum is free of negative-norm states. The latter is defined as the subspace of Virasoro primaries, satisfying the mass-shell condition, at criticality, namely when $c_{\text{int}} + \frac{3k}{k+2} = 26$ ($c_{\text{int}} > 0$, is the contribution of some internal CFT, and we are considering here the bosonic case). Within the present framework, the following conclusions can be drawn: (i) representations based on continuous series contain only positive-norm states, but they all describe tachyons; (ii) representations based on discrete series contain tachyons, massless and massive excitations, but for sufficiently large $|j|$ (j is negative so as to ensure unitarity of level-zero states), negative-norm states appear, which remain in the physical spectrum.

One possible solution to this problem is to cut the spin, hoping that all negative-norm states are thus eliminated [2, 3]. A refined analysis shows that a reasonable constraint is

$$\frac{k}{2} \leq j < 0.$$

Straightforward norm computations up to level two indicate that unitarity could be recovered [2], and there are even indications that a no-ghost theorem could exist [9]. However, an important issue is whether the above condition is consistent with other features of the string. The answer to this question is not known. Clearly such a constraint on the allowed values of the spin introduces a cut-off on the mass of string excitations. This is in obvious disagreement with the usual infinite tower of massive states, and, to some extent, with what we have learned so far from the ADS/CFT conjecture.

Another relevant and related question we must face concerns the one-loop partition function. Several problems appear there. In order to avoid the divergence due to the infinite degeneracy level by level, we must switch on an external source coupled to the Cartan generator. This allows for a proper – though not physically clear – definition of the characters of the discrete series [10]. It does not apply, however, successfully to continuous series, whose contribution to the partition function therefore remains undetermined. Ignoring that contribution, and concentrating on the discrete series, is not very conclusive either. On the one hand, one can seriously argue that, within the above unitarity constraint, modular invariance cannot be reached [11]. On the other hand,

even ignoring the unitarity bound, modular-invariant combinations of the characters are known when $k > -2$, but it is technically difficult to go beyond, since the characters themselves have not been computed, except for the case of irreducible representations.

As a conclusion, one can summarize the situation as follows: in the framework of the current-algebra approach, unitarity seems to be incompatible with a consistent physical spectrum, since it introduces a cut-off over the mass. This may be interpreted as a signal that, in the present approach, we are missing other sectors, which could provide us with positive-norm states, whose spin is $j < k/2$, and which might originate from a proper treatment of the *target-space boundary conditions*. The latter are hard to implement within the current-algebra method, but might be successfully explored following a path-integral approach *à la* Gawędzki [5]. The role of continuous series could also be clarified in this way. Although they are compatible with unitarity, the corresponding excitations seem to be all tachyonic. It is fair to mention here that one could try alternatively to avoid the mass/spin cut-off and the purely tachyonic continuous representations in various ways, playing essentially with the current algebra, and/or modifying the affine Sugawara construction [12]. However, it is not clear whether such modifications leave unaltered the interpretation of the theory as a string propagating over ADS_3 .

The motivations for studying the string on ADS_3 turn out to be wider than expected in the early works. The analysis of the ADS/CFT conjecture, in the framework of the $\text{ADS}_3 \times S^3$ background, in particular, has attracted most attention. Ideally, we would like to compute correlators in both sides and compare them. In practice, correlators for ADS_3 string states are out of reach, which makes any rigorous check quite intricate. Therefore, most of the work in that direction has been devoted to trying to express the space-time as well as the asymptotic two-dimensional conformal symmetry, in terms of the fields of the WZW model whose target space is the bulk ADS_3 theory, and to build in that way the boundary conformal field theory. It is fair to say, however, that this approach has not shed any light on the structure of the ADS_3 string itself – at least regarding the questions raised here; as long as the ADS_3 side is not handled exactly, the achievements are limited both on checking the ADS/CFT correspondence and on building the boundary theory [13]. Of course, there still is the – weaker – alternative to work with the low-energy supergravity, supplemented with all Kaluza–Klein excitations coming from higher dimensions, thus trying to obtain some feedback for the string on ADS_3 . For example, there are signs that all $SU(1,1)$ representations – discrete and continuous – should appear without bound on the mass.

If such a bound were present, assuming the ADS/CFT correspondence, it would be hard to identify states in the bulk ADS_3 supergravity (or string, as a fundamental theory), with states in the boundary conformal field theory.

Acknowledgments

It is a pleasure to thank C. Bachas and P. Bain for the stimulating collaboration we have been having on this subject. I would like to thank also the organizers of the Cargèse 99 ASI. The present note is a strict summary of a more extended version [11], where a complete bibliography can be found. This work was supported in part by EEC TMR contract ERBFMRX-CT96-0090.

References

- [1] J. Balog, L. O’Raifeartaigh, P. Forgács and A. Wipf, Nucl. Phys. **B325** (1989) 225.
- [2] P.M. Petropoulos, Phys. Lett. **236B** (1990) 151.
- [3] N. Mohameddi, Int. J. Mod. Phys. **A5** (1990) 3201.
- [4] I. Bars and D. Nemeschansky, Nucl. Phys. **B348** (1991) 89.
- [5] K. Gawędzki, proceedings of the Cargèse summer institute: “New symmetry principles in quantum field theory”, Cargèse, France, 16–27 July, 1991.
- [6] I. Antoniadis, C. Bachas and A. Sagnotti, Phys. Lett. **235B** (1990) 255.
- [7] M. Bañados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett. **69** (1992) 1849.
- [8] V. Balasubramanian, P. Kraus and A. Lawrence, Phys. Rev. **D59** (1999) 046003.
- [9] S. Hwang, Nucl. Phys. **B354** (1991) 100; M. Henningson and S. Hwang, Phys. Lett. **258B** (1991) 341; J. Evans, M. Gaberdiel and M. Perry, Nucl. Phys. **B535** (1998) 152.
- [10] V. Kač and M. Wakimoto, Proc. Nat. Acad. Sci. USA **85** (1988) 4956.
- [11] P.M. Petropoulos, hep-th/9908189.
- [12] I. Bars, Phys. Rev. **D53** (1996) 3308.
- [13] A. Giveon, D. Kutasov and N. Seiberg, Adv. Theor. Math. Phys. **2** (1998) 733; J. de Boer, H. Ooguri, H. Robins and J. Tannenhauser, JHEP **9812** (1998) 26; D. Kutasov and N. Seiberg, JHEP **9904** (1999) 8; A. Giveon and M. Roček, JHEP **9904** (1999) 19.